

Intern. Jour. of Diff. Eqs and Appl., 2, N3, (2001)

arXiv:math-ph/0102030v1 22 Feb 2001

New proof of Weyl's theorem ^{*†}

A.G. Ramm

Mathematics Department, Kansas State University,
Manhattan, KS 66506-2602, USA

ramm@math.ksu.edu

<http://www.math.ksu.edu/~ramm>

Abstract

Let $lu = -u'' + q(x)u$, where $q(x)$ is a real-valued $L^2_{loc}(0, \infty)$ function. H. Weyl has proved in 1910 that for any z , $Imz \neq 0$, the equation $(l - z)w = 0$, $x > 0$, has a solution $w \in L^2(0, \infty)$.

We prove this classical result using a new argument.

1 Introduction

Let $lu = -u'' + q(x)u$, where $q(x) \in L^2_{loc}$ is a real-valued function. Fix an arbitrary complex number z , $Imz > 0$, and consider the equation

$$lw - zw = 0, \quad x > 0 \tag{1.1}$$

H. Weyl proved [5] that equation (1.1) has a solution $w \in L^2(0, \infty)$, which is called a Weyl's solution. He gave the limit point-limit circle classification of the operator l : if equation (1.1) has only one solution $w \in L^2(0, \infty)$, then it is a limit point case, otherwise it is a limit circle case.

Weyl's theory is presented in several books, e.g. in [4], [3]. This theory is based on some limiting procedure $b \rightarrow \infty$ for the solutions to (1.1) on a finite interval $(0, b)$. In [3] a nice different proof is given for continuous $q(x)$.

The aim of our paper is to give a new method for a proof of Weyl's result.

Theorem 1.1. *Equation (1.1) has a solution $w \in L^2(0, \infty)$.*

*key words: limit circle, limit point, Weyl's solution.

†Math subject classification: 34B25, 34B20

Let us outline the new approach and the steps of the proof.

Since $q(x)$ is a real-valued function, symmetric operator l_0 defined on a linear dense subset $C_0^\infty(0, \infty)$ of $H = L^2(0, \infty)$ by the expression $lu = -u'' + q(x)u$ has a selfadjoint extension, which we denote by l . Therefore the resolvent $(l - z)^{-1}$ is a bounded linear operator on the Hilbert space

$$H = L^2(0, \infty), \quad \|(l - z)^{-1}\| \leq |Imz|^{-1}.$$

This operator is an integral operator with the kernel $G(x, y; z)$, which is a distribution satisfying the equation

$$(l - z)G(x, y; z) = \delta(x - y), \quad G(x, y; z) = G(y, x; z). \quad (1.2)$$

We will prove that

$$\int_0^\infty |G(x, y; z)|^2 dy \leq c(x; z) \quad \forall x \in (0, \infty), \quad Imz > 0, \quad (1.3)$$

where $c(x; z) = \text{const} > 0$.

The kernel $G(x, y; z)$, which is the Green function of the operator l , can be represented as

$$G(x, y; z) = \varphi(y; z)w(x; z), \quad x > y, \quad (1.4)$$

where w and φ are linearly independent solution to (1.1), so that $w(x; z) \not\equiv 0$. From (1.3) it follows that

$$w(x; z) \in L^2(0, \infty). \quad (1.5)$$

A detailed proof is given in section 2.

One may try to prove the existence of a Weyl's solution as follows: take an $h \in L_{loc}^2(0, \infty)$, $h = 0$ for $x > R$, $h \not\equiv 0$, and let $W := W(x, z) := (l - z)^{-1}h$, $Imz > 0$. Then W solves (1.1) for $x > R$ and $W \in L^2(0, \infty)$ since l is a selfadjoint operator in H . However, one has to prove then that W does not vanish identically for $x > R$, and this will be the case not for an arbitrary h with the above properties. In our paper the role of h is played by the delta-function, and since $\varphi(y; z)$ and w in (1.4) are linearly independent solutions of (1.1), one concludes that w does not vanish identically.

2 Proofs

Lemma 2.1. *If $q(x) \in L_{loc}^1(0, \infty)$ and $q(x)$ is real-valued, then symmetric operator*

$$l_0 u := -u'' + q(x)u, \quad D(l_0) = \{u : u \in C_0^\infty(0, \infty), \quad l_0 u \in H := L^2(0, \infty)\}$$

is defined on a linear dense in H subset, and admits a selfadjoint extension l .

Proof. This result is known: the density of the domain of definition of the symmetric operator l_0 mentioned in Lemma 1 and the existence of a selfadjoint extension are proved in [2]. The defect indices of l_0 are (1,1) or (2,2), so that by von Neumann extension theory l_0 has selfadjoint extensions (see [2]). Actually we assume in the Appendix that $q \in L^2_{loc}(0, \infty)$, in which case the conclusion of Lemma 2.1 is obvious: $C_0^\infty(0, \infty)$ is the linear dense subset in H on which l_0 is defined. \square

Let l be a selfadjoint extension of l_0 , $(l - z)^{-1}$ be its resolvent, $Imz > 0$, and $G(x, y; z)$ be the resolvent's kernel (in the sense of distribution theory) of $(l - z)^{-1}$, $G(x, y; z) = G(y, x; z)$.

Lemma 2.2. *For any fixed $x \in [0, \infty)$ one has*

$$\left(\int_0^\infty |G(x, y; z)|^2 dy \right)^{\frac{1}{2}} \leq c, \quad c = c(x; z) = \text{const} > 0. \quad (2.1)$$

Proof. Let $h \in C_0^\infty(0, \infty)$ and $u := (l - z)^{-1}h$, so

$$u(x; z) = \int_0^\infty G(x, y; z)h(y)dy, \quad (l - z)u = h. \quad (2.2)$$

Let us prove that:

$$|u(x; z)| \leq c(x; z)\|h\|, \quad (2.3)$$

where $x \in [0, \infty)$ is an arbitrary fixed point, $c(x) = \text{const} > 0$, $\|h\| := \|h\|_{L^2(0, \infty)}$, $(u, v) := (u, v)_{L^2(0, \infty)}$.

If (2.3) is proved, then

$$|(G(x, y; z), h)| \leq c(x; z)\|h\|. \quad (2.4)$$

From (2.4) the desired conclusion (2.1) follows immediately by the Riesz theorem about linear functionals in H .

To complete the proof, one has to prove estimate (2.3).

This estimate follows from the inequality:

$$\|u\|_{C(D_1)} \leq c (\| -u'' + q(x)u - zu \|_{L^2(D_2)} + \|u\|_{L^2(D_2)}) \leq c \left(1 + \frac{1}{|Imz|} \right) \|h\|, \quad (2.5)$$

where $c = c(D_1, D_2) = \text{const} > 0$, $D_1 \subset D_2$, $D_2 \subset [0, \infty)$, D_1 is a strictly inner open subinterval of D_2 .

Indeed, since l is selfadjoint, (2.2) implies:

$$\|u\| \leq \frac{\|h\|}{|Imz|}. \quad (2.6)$$

Moreover

$$-u'' + qu - zu = h, \quad (2.7)$$

so, using (2.6), one gets:

$$\|u\|_{L^2(D_2)} + \|-u'' + qu - zu\|_{L^2(D_2)} \leq \frac{\|h\|}{|Imz|} + \|h\| \leq \left(1 + \frac{1}{|Imz|}\right) \|h\|, \quad (2.8)$$

From (2.5), (2.6) and (2.8) one gets (2.3).

Let us finish the proof by proving (2.5).

In fact, inequality (2.5) is a particular case of the well-known elliptic estimates (see e.g. [1, pp. 239-241]), but an elementary proof of (2.5) is given below in the Appendix.

Lemma 2 is proved. \square

Proof of Theorem 1.1

Equation (1.2) implies that

$$G(x, y; z) = \varphi(x; z)w(y; z), \quad y \geq x,$$

where $w(y; z)$ solves (1.1), and the function $\varphi(x; z)$ is also a solution to (1.1). Inequality (2.1) implies $w \in L^2(0, \infty)$ if $Imz > 0$.

Theorem 1.1 is proved. \square

To make this paper self-contained we give an elementary proof of inequality (2.5) in the Appendix. This proof allows one to avoid reference to the elliptic inequalities [1], the proof of which in [1] is long and complicated (in [1] the multidimensional elliptic equations of general form are studied, which is the reason for the complicated argument in [1]).

Appendix: An elementary proof of inequality (2.5).

Since $u(x)$ is $C_{loc}^1(0, \infty)$ it is sufficient to prove (2.5) assuming that $D_1 = (a, b)$ and $b - a$ is arbitrarily small. Let $\eta(x) \in C_0^\infty(a, b)$ be a cut-off function, $0 \leq \eta \leq 1$, $\eta(x) = 1$ in $(a + \delta, b - \delta)$, $0 < \delta < \frac{b-a}{4}$, $\eta(x) = 0$ in a neighborhoods of points a and b .

Let $v = \eta u$. Then (2.2) implies:

$$lv = \eta h - 2\eta' u' - \eta'' u, \quad v(a) = v'(a) = 0.$$

Thus

$$v'' = qv - zv - \eta h + \eta'' u + 2\eta' u', \quad (A.1)$$

and

$$\begin{aligned} |v(x)| &= \left| \int_a^x (x-s)v''(s)ds \right| \leq c_1 \int_a^b [|qv| + |z||v|] ds + c_2, \\ &\int_a^b |h|ds + c_2 \int_a^b |u|ds + c_2 \int_a^b |u'|ds. \end{aligned} \quad (A.2)$$

Here

$$c_1 = b - a, c_2 = \max_{a \leq x \leq b} [|\eta(x)| + |\eta'' + 2|\eta'|].$$

If $b - a$ is sufficiently small, then

$$c_1 \int_a^b (|q| + |z|) dx \max_{a \leq x \leq b} |v(x)| < \gamma \max_{a \leq x \leq b} |v(x)|, \quad 0 < \gamma < 1.$$

Therefore (A.1) implies

$$\max_{a \leq x \leq b} |v(x)| \leq c_3 [\|h\|_{L^2(a,b)} + \|u\|_{L^2(a,b)} = \|u'\|_{L^2(a,b)}], \quad (\text{A.3})$$

where $c_3 = c_3(a, b; z)$. From (A.3) and (2.6) it follows that inequality (2.5) holds, provided that:

$$\|u'\|_{L^2(a,b)} \leq c\|h\| + \delta\|u\|_{L^\infty}. \quad (\text{A.4})$$

The last estimate is proved as follows. Multiply (2.2) by $\eta\bar{u}$ (the bar stands for complex conjugate and η is a cut-off function, $\eta \in C_0^\infty(a, b)$ and integrate over (a, b) to get

$$\int_a^b |u'|^2 \eta dx = \int_a^b u' \bar{u} \eta' dx + \int_a^b \eta h \bar{u} dx + z \int_a^b \eta |u|^2 dx - \int_a^b q |u|^2 \eta dx := I_1 + I_2 + I_3 + I_4.$$

One has, using the inequality $|uv| \leq \varepsilon|u|^2 + \frac{|v|^2}{4\varepsilon}$, $\varepsilon > 0$,

$$|I_1| \leq c \left(\varepsilon \|u'\|^2 + \frac{1}{4\varepsilon} \|u\|^2 \right), \quad c = \max |\eta'|,$$

$$|I_2| + |I_3| \leq c (\|h\| \|u\| + \|u\|^2) \leq c_1 \|h\|^2,$$

where (2.6) was used,

$$|I_4| \leq \|qu\| \|u\| \leq \|q\|_{L^2} \|u\|_{L^\infty} \|u\|.$$

Thus, if $a < a_1 < b_1 < b$, where $\eta = 1$ on $[a_1, b_1]$, one gets

$$\int_{a_1}^{b_1} |u'|^2 dx \leq C (\|h\|^2 + \|u\|_{L^\infty} \|h\|) \leq \delta \|u\|_{L^\infty}^2 + C \|h\|^2, \quad (\text{A.5})$$

where $C = C(\varepsilon, z, a, b, \delta) = \text{const} > 0$, $0 < \delta$ can be chosen arbitrarily small. Inequality (A.5) implies (A.4).

Inequality (2.5) is proved. \square

References

- [1] Gilbarg, D., Trudinger, N., Elliptic partial differential equations of second order, Springer, New York, 1983
- [2] Naimark, M., Linear differential operators, Ungar, New York, 1968.
- [3] Reed, M., Simon, B., Methods of modern mathematical physics, vol.2, Acad. Press, New York, 1978.
- [4] Titchmarsh E., Eigenfunction expansions associated with second-order differential equations, Oxford, Clarendon Press, 1946.
- [5] Weyl, H., Über gewöhnliche Differentialgleichungen mit Singularitäten and die zugehörigen Entwicklungen willkürliche Funktionen, Math. Ann., 68, (1910), 220-269.